arXiv:1611.03302v1 [math.GR] 10 Nov 2016

The number of subgroups of the group $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s$

László Tóth

Abstract

We deduce direct formulas for the total number of subgroups and the number of subgroups of a given order of the group $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s$, where $m, n, r, s \in \mathbb{N}$. The proofs are by some simple group theoretical and number theoretical arguments based on Goursat's lemma for groups. Two conjectures are also formulated.

2010 Mathematics Subject Classification: 20K01, 20K27, 11A25, 11Y70

Key Words and Phrases: cyclic group, direct product, subgroup, number of subgroups, Goursat's lemma, finite abelian group of rank four

1 Introduction

Throughout the paper we use the following notation: $\mathbb{N} := \{1, 2, \ldots\}$, $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$; $\mathbb{P}$ is the set of primes; the prime power factorization of $n \in \mathbb{N}$ is $n = \prod_{p\in\mathbb{P}} p^{\nu_p(n)}$, where all but a finite number of the exponents $\nu_p(n)$ are zero; $\mathbb{Z}_n$ denotes the additive group of residue classes modulo n ($n \in \mathbb{N}$); $\gcd(a, b)$ and $\operatorname{lcm}(a, b)$ denote the greatest common divisor and least common multiple, respectively of $a, b \in \mathbb{N}$; φ is Euler's arithmetic function.

For an arbitrary finite abelian group G of order n let $N(G)$ and $N(G; k)$ denote the total number of subgroups, respectively the number of subgroups of order k of G ($k \mid n$). Let $G = \times_{p\in\mathbb{P}} G_p$ be the primary decomposition of G, where $|G_p| = p^{\nu_p(n)}$ ($p \in \mathbb{P}$). Then

$$N(G) = \prod_{p\in\mathbb{P}} N(G_p), \tag{1}$$

$$N(G; k) = \prod_{p\in\mathbb{P}} N(G_p; p^{\nu_p(k)}). \tag{2}$$

Therefore, the problem of counting the subgroups of G reduces to p-groups. It is known that the number of subgroups of a given type in an abelian p-group is a polynomial in p with positive integral coefficients, which can be represented in terms of the gaussian coefficients. There are also various other formulas in the literature. We refer to the monograph by Butler [4] and the papers by Bhowmik [2], Călugăreanu [5], Shokuev [13], Stehling [14].

It follows that in an abelian p-group the total number of subgroups and the number of subgroups of a given order are also polynomials in p with positive integral coefficients. However,

it is a difficult task to make explicit these polynomials, even in the case of p-groups of small rank. The rank two polynomials are

$$N(\mathbb{Z}_{p^a} \times \mathbb{Z}_{p^b}) = \frac{(b-a+1)p^{a+2} - (b-a-1)p^{a+1} - (a+b+3)p + (a+b+1)}{(p-1)^2}, \tag{3}$$

and

$$N(\mathbb{Z}_{p^a} \times \mathbb{Z}_{p^b}; p^c) = \begin{cases} \frac{p^{c+1}-1}{p-1}, & c \leqslant a \leqslant b, \\ \frac{p^{a+1}-1}{p-1}, & a \leqslant c \leqslant b, \\ \frac{p^{a+b-c+1}-1}{p-1}, & a \leqslant b \leqslant c \leqslant a+b, \end{cases} \tag{4}$$

representing the total number of subgroups, respectively the number of subgroups of order p^c of the rank two group $\mathbb{Z}_{p^a} \times \mathbb{Z}_{p^b}$ with $1 \leqslant a \leqslant b$, $0 \leqslant c \leqslant a+b$. See Călugăreanu [5], Petrillo [12, Prop. 2], Tărnăuceanu [16, Th. 3.3]).

Now consider the group $\mathbb{Z}_m \times \mathbb{Z}_n$, where $m, n \in \mathbb{N}$. Note that $\mathbb{Z}_m \times \mathbb{Z}_n$ is isomorphic to $\mathbb{Z}_{\gcd(m,n)} \times \mathbb{Z}_{\mathrm{lcm}(m,n)}$, and it has rank two, assuming that $\gcd(m,n) > 1$. Recall that a finite abelian group of order > 1 has rank t if it is isomorphic to $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_t}$, where $n_1, \ldots, n_t \in \mathbb{N}\backslash\{1\}$ and $n_j \mid n_{j+1}$ $(1 \leqslant j \leqslant t-1)$, which is the invariant factor decomposition of the given group. Here the number t is uniquely determined and represents the minimal number of generators of the given group.

For every $m, n \in \mathbb{N}$ the number of all subgroups of $\mathbb{Z}_m \times \mathbb{Z}_n$ is

$$N(\mathbb{Z}_m \times \mathbb{Z}_n) = \sum_{\substack{i \mid m \\ j \mid n}} \gcd(i,j), \tag{5}$$

and the number of its subgroups of order k $(k \mid mn)$ is given by

$$N(\mathbb{Z}_m \times \mathbb{Z}_n; k) = \sum_{\substack{i \mid \gcd(m,k) \\ j \mid \gcd(n,k) \\ k \mid ij}} \varphi\left(\frac{ij}{k}\right). \tag{6}$$

Formulas (5) and (6), established by the author [18, Ths. 4.1, 4.3], generalize and put in more compact forms identities (3) and (4), respectively. See also the paper by Hampejs, Holighaus, the author, and Wiesmeyr [8] for a different approach.

The polynomial giving the total number of subgroups of the rank three group $\mathbb{Z}_{p^a} \times \mathbb{Z}_{p^b} \times \mathbb{Z}_{p^c}$ was obtained by Oh [11, Cor. 2.2] and by Tărnăuceanu and the author [17, Cor. 2.9]. Formulas similar to (5) and (6), concerning the numbers of all subgroups, respectively subgroups of order k of the group $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r$ $(m, n, r, k \in \mathbb{N}, k \mid mnr)$ were given by Hampejs and the author [9, Th. 2.2].

The next step is to investigate rank four groups, which is the goal of the present paper. The special case of the group $\mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_{p^i}$ $(i \in \mathbb{N})$ was recently considered by Chew, Chin and Lim [6]. Instead of p-groups we prefer to investigate the group $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s$ with $m, n, r, s \in \mathbb{N}$, which has rank four, provided that $\gcd(m,n,r,s) > 1$. We deduce direct formulas for

$$N(m,n,r,s) := N(\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s)$$

and

$$N(m,n,r,s;k) := N(\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s; k),$$

representing the total number of subgroups of the group $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s$ and the number of its subgroups of order k $(m,n,r,s,k \in \mathbb{N}, k \mid mnrs)$. See Theorems 5 and 6. As direct consequences, in Corollaries 7 and 8 we derive formulas for the number of subgroups of p-groups of rank four. We do not obtain the explicit form of the corresponding polynomials, but those can be computed in special cases. In Section 5 we present tables for such polynomials, computed using Corollaries 7 and 8. We also formulate two conjectures.

For the proofs of our results we use some simple group theoretical and number theoretical arguments based on Goursat's lemma for groups (Proposition 1). We remark that for every $m,n,r,s \in \mathbb{N}$,

$$N(m,n,r,s) = \prod_{p\in\mathbb{P}} N(p^{\nu_p(m)}, p^{\nu_p(n)}, p^{\nu_p(r)}, p^{\nu_p(s)}), \tag{7}$$

$$N(m,n,r,s;k) = \prod_{p\in\mathbb{P}} N(p^{\nu_p(m)}, p^{\nu_p(n)}, p^{\nu_p(r)}, p^{\nu_p(s)}; p^{\nu_p(k)}),$$

which are direct consequences of the identities (1) and (2). Here (7) shows that $(m,n,r,s) \mapsto N(m,n,r,s)$ is a multiplicative arithmetic function of four variables. We refer to our survey paper [19] regarding this number theoretical concept.

In turns out that the function $n \mapsto N(n) := N(n,n,n,n)$ is multiplicative, that is $N(n) = \prod_{p\in\mathbb{P}} N(p^{\nu_p(n)})$ for every $n \in \mathbb{N}$. A table of values for $N(n)$ is presented in Section 5.

2 Preliminaries

Goursat's lemma for groups can be stated as follows. For its history, proof and discussion see the references given in [18]. See also the recent paper [1].

Proposition 1 (Goursat's lemma for groups)**.** *Let G and H be arbitrary groups. Then there is a bijection between the set S of all subgroups of $G\times H$ and the set T of all 5-tuples (A,B,C,D,Ψ), where $B \trianglelefteq A \leqslant G$, $D \trianglelefteq C \leqslant H$ and $\Psi : A/B \to C/D$ is an isomorphism (here $\leqslant$ denotes subgroup and $\trianglelefteq$ denotes normal subgroup). More precisely, the subgroup corresponding to (A,B,C,D,Ψ) is*

$$K = \{(g,h) \in A \times C : \Psi(gB) = hD\}.$$

We will need the following corollary.

Corollary 2. *Assume that G and H are finite groups and that the subgroup K of $G \times H$ corresponds to the 5-tuple $(A_K, B_K, C_K, D_K, \Psi_K)$ under this bijection. Then one has $|A_K| \cdot |D_K| = |K| = |B_K| \cdot |C_K|$.*

Proposition 1 was used in our paper [18] to count the subgroups of the group $(\mathbb{Z}_m \times \mathbb{Z}_n, +)$ $(m,n \in \mathbb{N})$. In fact, formulas (5) and (6) are consequences of the next representation result. For every $m,n \in \mathbb{N}$ let

$$J_{m,n} := \left\{(a,b,c,d,\ell) \in \mathbb{N}^5 : a \mid m, b \mid a, c \mid n, d \mid c, \frac{a}{b} = \frac{c}{d}, \ell \leqslant \frac{a}{b}, \gcd\left(\ell, \frac{a}{b}\right) = 1\right\}. \tag{8}$$

Note that here $\gcd(b,d)\cdot\operatorname{lcm}(a,c)=ad$ and $\gcd(b,d)\mid\operatorname{lcm}(a,c)$.

For $(a,b,c,d,\ell)\in J_{m,n}$ define

$$K_{a,b,c,d,\ell}:=\left\{\left(i\frac{m}{a},i\ell\frac{n}{c}+j\frac{n}{d}\right):0\leqslant i\leqslant a-1,0\leqslant j\leqslant d-1\right\}.$$

Proposition 3 ([18, Th. 3.1])**.** *Let $m,n\in\mathbb{N}$.*

i) The map $(a,b,c,d,\ell)\mapsto K_{a,b,c,d,\ell}$ is a bijection between the set $J_{m,n}$ and the set of subgroups of $(\mathbb{Z}_m\times\mathbb{Z}_n,+)$.

ii) The invariant factor decomposition of the subgroup $K_{a,b,c,d,\ell}$ of order ad is

$$K_{a,b,c,d,\ell}\simeq\mathbb{Z}_{\gcd(b,d)}\times\mathbb{Z}_{\operatorname{lcm}(a,c)}.$$

We need the following new result concerning the quotient group $(\mathbb{Z}_m\times\mathbb{Z}_n)/K_{a,b,c,d,\ell}$. Its proof is included in Section 4.

Proposition 4. *Let $m,n\in\mathbb{N}$. For every subgroup $K_{a,b,c,d,\ell}$ defied above, we have the invariant factor decomposition*

$$(\mathbb{Z}_m\times\mathbb{Z}_n)/K_{a,b,c,d,\ell}\simeq\mathbb{Z}_{\gcd\left(\frac{m}{a},\frac{n}{c}\right)}\times\mathbb{Z}_{\operatorname{lcm}\left(\frac{m}{b},\frac{n}{d}\right)},$$

where $\gcd\left(\frac{m}{a},\frac{n}{c}\right)\mid\operatorname{lcm}\left(\frac{m}{b},\frac{n}{d}\right)$.

Let $F(m,n)$ denote the number of automorphisms of the group $\mathbb{Z}_m\times\mathbb{Z}_n$. For every $m,n\in\mathbb{N}$,

$$F(m,n)=\prod_{p\in\mathbb{P}}F(p^{\nu_p(m)},p^{\nu_p(n)}),$$

hence $(m,n)\mapsto F(m,n)$ is a multiplicative arithmetic function of two variables. For prime powers p^a,p^b one has, cf. [10, Th. 4.1], [7, Cor. 3],

$$F(p^a,p^b)=\begin{cases}p^{2a}\varphi(p^a)\varphi(p^b), & 0\leqslant a<b,\\ p^a\varphi_2(p^a)\varphi(p^a), & 0\leqslant a=b,\end{cases}$$

where φ_2 is the Jordan function of order 2 given by $\varphi_2(n)=n^2\prod_{p\mid n}(1-\frac{1}{p^2})$. We deduce that $F(1,p^b)=\varphi(p^b)$ for $b\in\mathbb{N}$ and

$$F(p^a,p^b)=\begin{cases}p^{3a+b}\left(1-\frac{1}{p}\right)^2, & 1\leqslant a<b,\\ p^{4a}\left(1-\frac{1}{p}\right)\left(1-\frac{1}{p^2}\right), & 1\leqslant a=b.\end{cases}$$

3 Main results

We state our main results. Their proofs are included in Section 4.

Theorem 5. *For every $m, n, r, s \in \mathbb{N}$ we have*

$$N(m,n,r,s) = \sum \varphi(x_3)\varphi(y_3)\varphi(z_3)\varphi(t_3)F(u,v), \tag{9}$$

where the sum is over all $(x_1, x_2, x_3, x_4, x_5, y_1, y_2, y_3, y_4, y_5, z_1, z_2, z_3, z_4, z_5, t_1, t_2, t_3, t_4, t_5) \in \mathbb{N}^{20}$ satisfying the following 10 conditions:

(1) $x_1x_2x_3 = m$,
(2) $x_3x_4x_5 = n$,
(3) $y_1y_2y_3 = \gcd(x_1, x_4)$,
(4) $y_3y_4y_5 = x_3 \operatorname{lcm}(x_1, x_4)$,
(5) $z_1z_2z_3 = r$,
(6) $z_3z_4z_5 = s$,
(7) $t_1t_2t_3 = \gcd(z_1, z_4)$,
(8) $t_3t_4t_5 = z_3 \operatorname{lcm}(z_1, z_4)$,
(9) $\gcd(y_2, y_5) = \gcd(t_2, t_5) =: u$,
(10) $y_3 \operatorname{lcm}(y_2, y_5) = t_3 \operatorname{lcm}(t_2, t_5) =: v$.

Theorem 6. *For every $k, m, n, r, s \in \mathbb{N}$ such that $k \mid mnrs$, the number $N(m,n,r,s;k)$ is given by the sum (9) with the additional constraint*

(11) $x_1x_3x_4t_1t_3t_4 = k$.

Corollary 7. *For every $a, b, c, d \in \mathbb{N}_0$ we have*

$$N(p^a, p^b, p^c, p^d) = \sum \varphi(p^{x_3})\varphi(p^{y_3})\varphi(p^{z_3})\varphi(p^{t_3})F(p^u, p^v), \tag{10}$$

where the sum is over all $(x_1, x_2, x_3, x_4, x_5, y_1, y_2, y_3, y_4, y_5, z_1, z_2, z_3, z_4, z_5, t_1, t_2, t_3, t_4, t_5) \in \mathbb{N}_0^{20}$ satisfying the following 10 conditions:

(i) $x_1 + x_2 + x_3 = a$,
(ii) $x_3 + x_4 + x_5 = b$,
(iii) $y_1 + y_2 + y_3 = \min(x_1, x_4)$,
(iv) $y_3 + y_4 + y_5 = x_3 + \max(x_1, x_4)$,
(v) $z_1 + z_2 + z_3 = c$,
(vi) $z_3 + z_4 + z_5 = d$,
(vii) $t_1 + t_2 + t_3 = \min(z_1, z_4)$,
(viii) $t_3 + t_4 + t_5 = z_3 + \max(z_1, z_4)$,
(ix) $\min(y_2, y_5) = \min(t_2, t_5) =: u$,
(x) $y_3 + \max(y_2, y_5) = t_3 + \max(t_2, t_5) =: v$.

Corollary 8. *For every $a, b, c, d, k \in \mathbb{N}_0$ with $k \leqslant a + b + c + d$ the number $N(p^a, p^b, p^c, p^d; p^k)$ is given by the sum (10) with the additional constraint*

(xi) $x_1 + x_3 + x_4 + t_1 + t_3 + t_4 = k$.

4 Proofs

Proof of Proposition 4. Let $K := K_{a,b,c,d,\ell}$. The group $\mathbb{Z}_m \times \mathbb{Z}_n$ is generated by $(0,1)$ and $(1,0)$. Therefore, the quotient group $\mathbb{Z}_m \times \mathbb{Z}_n/K$ is generated by $(0,1)+K$ and $(1,0)+K$.

First we show that the order of $(0,1)+K$ is $\frac{n}{d}$. Indeed, this follows from the following properties:

- $t(0,1) \in K$ if and only if there is $(i,j) \in \{0,1,\dots,a-1\} \times \{0,1,\dots,d-1\}$ such that $(0,t) = (i\frac{m}{a}, i\ell\frac{n}{c} + j\frac{n}{d})$, where the last condition is equivalent, in turn, to $i=0$ and $t = j\frac{n}{d}$;
- the least such $t \in \mathbb{N}$ is $\frac{n}{d}$.

Next we show that the order of $(1,0)+K$ is $\frac{m}{b}$. Indeed, observe that

- $u(1,0) \in K$ if and only if there is $(i,j) \in \{0,1,\dots,a-1\} \times \{0,1,\dots,d-1\}$ such that $(u,0) = (i\frac{m}{a}, i\ell\frac{n}{c} + j\frac{n}{d})$;
- therefore, $u = i\frac{m}{a}$ and $i\ell\frac{n}{c} + j\frac{n}{d} \equiv 0 \pmod{n}$, that is $\frac{n}{c}(i\ell + j\frac{c}{d}) \equiv 0 \pmod{n}$, $i\ell + j\frac{c}{d} \equiv 0 \pmod{c}$, the latter is a linear congruence in j and it turns out that for a fixed i it has solution in j if $\gcd\left(\frac{c}{d}, c\right) = \frac{c}{d} \mid i\ell$, that is $\frac{c}{d} \mid i$, since $\gcd\left(\ell, \frac{c}{d}\right) = 1$, cf. (8);
- the least such $i \in \mathbb{N}$ is $\frac{c}{d}$;
- finally, $u = \frac{c}{d} \cdot \frac{m}{a} = \frac{a}{b} \cdot \frac{m}{a} = \frac{m}{b}$, representing the order of $(1,0)+K$.

We deduce that the exponent of $(\mathbb{Z}_m \times \mathbb{Z}_n)/K$ is $\operatorname{lcm}(\frac{m}{b}, \frac{n}{d})$.

Now, $\mathbb{Z}_m \times \mathbb{Z}_n \simeq \mathbb{Z}_{\gcd(m,n)} \times \mathbb{Z}_{\operatorname{lcm}(m,n)}$ is a group of rank $\leqslant 2$. Therefore, the quotient group $(\mathbb{Z}_m \times \mathbb{Z}_n)/K$ has also rank $\leqslant 2$. That is, $(\mathbb{Z}_m \times \mathbb{Z}_n)/K \simeq \mathbb{Z}_v \times \mathbb{Z}_w$ for unique v and w, where $v \mid w$ and $vw = \frac{mn}{ad}$. We obtain that the exponent of $(\mathbb{Z}_m \times \mathbb{Z}_n)/K$ is $\operatorname{lcm}(\frac{m}{b}, \frac{n}{d}) = w$. Hence

$$v = \frac{mn}{adw} = \frac{mn}{ad\operatorname{lcm}(\frac{m}{b}, \frac{n}{d})} = \frac{mn}{ad\operatorname{lcm}(\frac{mn}{bn}, \frac{mn}{dm})} = \frac{mn}{ad\frac{mn}{\gcd(bn,dm)}}$$

$$= \frac{\gcd(bn,dm)}{ad} = \gcd\left(\frac{n}{d}\cdot\frac{b}{a}, \frac{m}{a}\right) = \gcd\left(\frac{n}{d}\cdot\frac{d}{c}, \frac{m}{a}\right) = \gcd\left(\frac{m}{a}, \frac{n}{c}\right).$$

This completes the proof. □

Proof of Theorem 5. We apply Goursat's lemma (Proposition 1) for the groups $G = \mathbb{Z}_m \times \mathbb{Z}_n$ and $H = \mathbb{Z}_r \times \mathbb{Z}_s$. Let

$$A = K_{a,b,c,d,\ell} \leqslant \mathbb{Z}_m \times \mathbb{Z}_n,$$

where $a \mid m, b \mid a, c \mid n, d \mid c, \frac{a}{b} = \frac{c}{d} = e, 1 \leqslant \ell \leqslant e, \gcd(\ell, e) = 1$. Here, by Proposition 3,

$$A \simeq \mathbb{Z}_{m_1} \times \mathbb{Z}_{n_1},$$

where $m_1 = \gcd(b,d)$, $n_1 = \operatorname{lcm}(a,c)$. Also, let

$$B \simeq K_{a_1,b_1,c_1,d_1,\ell_1} \leqslant A,$$

where $a_1 \mid m_1, b_1 \mid a_1, c_1 \mid n_1, d_1 \mid c_1, \frac{a_1}{b_1} = \frac{c_1}{d_1} = e_1, 1 \leqslant \ell_1 \leqslant e_1, \gcd(\ell_1, e_1) = 1$. We have, by using Proposition 4,

$$A/B \simeq \mathbb{Z}_u \times \mathbb{Z}_v, \tag{11}$$

where $u = \gcd(m_1/a_1, n_1/c_1)$, $v = \operatorname{lcm}(m_1/b_1, n_1/d_1)$.

In a similar way, let

$$C = K_{\alpha,\beta,\gamma,\delta,\lambda} \leqslant \mathbb{Z}_r \times \mathbb{Z}_s,$$

where $\alpha \mid r, \beta \mid \alpha, \gamma \mid s, \delta \mid \gamma, \frac{\alpha}{\beta} = \frac{\gamma}{\delta} = \epsilon, 1 \leqslant \lambda \leqslant \epsilon, \gcd(\lambda, \epsilon) = 1$. Here,

$$C \simeq \mathbb{Z}_{r_1} \times \mathbb{Z}_{s_1},$$

where $r_1 = \gcd(\beta, \delta)$, $s_1 = \operatorname{lcm}(\alpha, \gamma)$. Furthermore, let

$$D \simeq K_{\alpha_1,\beta_1,\gamma_1,\delta_1,\lambda_1} \leqslant C,$$

with $\alpha_1 \mid r_1, \beta_1 \mid \alpha_1, \gamma_1 \mid s_1, \delta_1 \mid \gamma_1, \frac{\alpha_1}{\beta_1} = \frac{\gamma_1}{\delta_1} = \epsilon_1, 1 \leqslant \lambda_1 \leqslant \epsilon_1, \gcd(\lambda_1, \epsilon_1) = 1$. Here,

$$C/D \simeq \mathbb{Z}_\eta \times \mathbb{Z}_\theta, \tag{12}$$

where $\eta = \gcd(r_1/\alpha_1, s_1/\gamma_1)$, $\theta = \operatorname{lcm}(r_1/\beta_1, s_1/\delta_1)$.

Assume that the quotient groups A/B and C/D are isomorphic. According to (11) and (12), this holds if and only if $u = \eta$ and $v = \theta$. Then there are $F(u, v)$ isomorphisms $A/B \simeq C/D$.

It follows that

$$N(m, n, r, s) = \sum F(u, v),$$

where the sum is over all $(a, b, c, d, \ell, a_1, b_1, c_1, d_1, \ell_1, \alpha, \beta, \gamma, \delta, \lambda, \alpha_1, \beta_1, \gamma_1, \delta_1, \lambda_1) \in \mathbb{N}^{20}$ satisfying the above properties.

More exactly, let $m = ax, a = by, n = cz, c = dt$, where $y = t = e$. Hence $m = bxe$, $n = dze$. Also, $m_1 = b_1x_1e_1$, $n_1 = d_1z_1e_1$, where $m_1 = \gcd(b, d)$, $n_1 = \operatorname{lcm}(a, c) = e \operatorname{lcm}(b, d)$. Similarly, let $r = \alpha X, \alpha = \beta Y$, $s = \gamma Z, \gamma = \delta T$, where $Y = T = \epsilon$. Hence $r = \beta X\epsilon$, $s = \delta Z\epsilon$. Also, $r_1 = \beta_1X_1\epsilon_1$, $s_1 = \delta_1Z_1\epsilon_1$, where $r_1 = \gcd(\beta, \delta)$, $r_1 = \operatorname{lcm}(\alpha, \gamma) = \epsilon \operatorname{lcm}(\beta, \delta)$.

We deduce that

$$N(m, n, r, s) = \sum F(u, v),$$

where the sum is over all $(b, x, e, d, z, \ell, b_1, x_1, e_1, d_1, z_1, \ell_1, \beta, X, \epsilon, \delta, Z, \lambda, \beta_1, X_1, \epsilon_1, \delta_1, Z_1, \lambda_1) \in \mathbb{N}^{24}$ satisfying the following conditions:

(1') $bxe = m$,
(2') $dze = n$,
(3') $b_1x_1e_1 = \gcd(b, d)$,
(4') $d_1z_1e_1 = e \operatorname{lcm}(b, d)$,
(4") $1 \leqslant \ell \leqslant e, \gcd(\ell, e) = 1$,
(5') $\beta X\epsilon = r$,
(6') $\delta Z\epsilon = s$,
(7') $\beta_1X_1\epsilon_1 = \gcd(\beta, \delta)$,
(8') $\delta_1Z_1\epsilon_1 = \epsilon \operatorname{lcm}(\beta, \delta)$,
(8") $1 \leqslant \ell_1 \leqslant e_1, \gcd(\ell_1, e_1) = 1$,
(9') $\gcd(x_1, z_1) = \gcd(X_1, Z_1) = u$,
(10') $e_1 \operatorname{lcm}(x_1, z_1) = \epsilon_1 \operatorname{lcm}(X_1, Z_1) = v$,
(10") $1 \leqslant \lambda \leqslant \epsilon, \gcd(\lambda, \epsilon) = 1$,
(10"') $1 \leqslant \lambda_1 \leqslant \epsilon_1, \gcd(\lambda_1, \epsilon_1) = 1$.

Note that, according to (4"), (8"), (10") and (10"'), the variables ℓ, ℓ_1, λ and λ_1 take $\varphi(e)$, $\varphi(e_1)$, $\varphi(\epsilon)$, respectively $\varphi(\epsilon_1)$ distinct values, independently from the other variables.

Now formula (9) follows by introducing the notation:

$x_1 := b$, $x_2 := x$, $x_3 := e$, $x_4 := d$, $x_5 := z$, $y_1 := b_1$, $y_2 := x_1$, $y_3 := e_1$, $y_4 := d_1$, $y_5 := z_1$ $z_1 := \beta$, $z_2 := X$, $z_3 := \epsilon$, $z_4 := \delta$, $z_5 := Z$, $t_1 := \beta_1$, $t_2 := X_1$, $t_3 := \epsilon_1$, $t_4 := \delta_1$, $t_5 := Z_1$. □

Proof of Theorem 6. According to Corollary 2, $k = |A||D| = |B||C|$. Here $|A| = ad = bde$, $|B| = a_1d_1 = b_1d_1e_1$. Similarly, $|C| = \alpha\delta = \beta\delta\epsilon$, $|D| = \alpha_1\delta_1 = \beta_1\delta_1\epsilon_1$. Hence condition $k = |A||D| = |B||C|$ is equivalent to $bde\beta_1\delta_1\epsilon_1 = k$ and to $x_1x_3x_4t_1t_3t_4 = k$ with the changing of notation of above. Note that relation $k = |B||C|$ is a consequence of the other constraints and needs not to be considered. □

Proof of Corollary 7. Apply Theorem 5 in the case $(m, n, r, s) = (p^a, p^b, p^c, p^d)$. □

Proof of Corollary 8. Apply Theorem 6 in the case $(m, n, r, s) = (p^a, p^b, p^c, p^d)$, $k := p^k$. □

5 Tables and conjectures

5.1 Values of $N(p^a, p^b, p^c, p^d)$ and $N(p^a, p^b, p^c, p^d; p^k)$

Consider the polynomials $N(p^a, p^b, p^c, p^d)$ and $N(p^a, p^b, p^c, p^d; p^k)$, where p is prime, $1 \leqslant a \leqslant b \leqslant c \leqslant d$, $0 \leqslant k \leqslant a+b+c+d =: n$. It is known that $N(p^a, p^b, p^c, p^d; p^k) = N(p^a, p^b, p^c, p^d; p^{n-k})$ for every $0 \leqslant k \leqslant n$. The following unimodality result is also known: $N(p^a, p^b, p^c, p^d; p^k) - N(p^a, p^b, p^c, p^d; p^{k-1})$ has nonnegative coefficients for every $1 \leqslant k \leqslant \lfloor n/2 \rfloor$. More generally, similar properties hold for every abelian p-group. See Butler [3], Takegahara [15].

In what follows we give the polynomials representing the values of $N(p^a, p^b, p^c, p^d)$ and $N(p^a, p^b, p^c, p^d; p^k)$, where $1 \leqslant a \leqslant b \leqslant c \leqslant d \leqslant 3$, also $a = b = c = d = 4$, with $0 \leqslant k \leqslant \lfloor n/2 \rfloor$, computed using Corollaries 7 and 8.

We also formulate the following conjectures, confirmed by these examples.

Conjecture 9. *For every $1 \leqslant a \leqslant b \leqslant c \leqslant d$, the degree of the polynomial $N(p^a, p^b, p^c, p^d)$ is $2a + b + c$.*

Conjecture 10. *For every $1 \leqslant m$ the degree of the polynomial $N(p^m, p^m, p^m, p^m)$ is $4m$ and its leading coefficient is 1.*

$N(p, p, p, p)$	$5 + 3p + 4p^2 + 3p^3 + p^4$
$N(p, p, p, p; 1)$	1
$N(p, p, p, p; p)$	$1 + p + p^2 + p^3$
$N(p, p, p, p; p^2)$	$1 + p + 2p^2 + p^3 + p^4$

$N(p, p, p, p^2)$	$6 + 4p + 6p^2 + 6p^3 + 2p^4$
$N(p, p, p, p^2; 1)$	1
$N(p, p, p, p^2; p)$	$1 + p + p^2 + p^3$
$N(p, p, p, p^2; p^2)$	$1 + p + 2p^2 + 2p^3 + p^4$

$N(p,p,p,p^3)$	$7+5p+8p^2+9p^3+3p^4$
$N(p,p,p,p^3;1)$	1
$N(p,p,p,p^3;p)$	$1+p+p^2+p^3$
$N(p,p,p,p^3;p^2)$	$1+p+2p^2+2p^3+p^4$
$N(p,p,p,p^3;p^3)$	$1+p+2p^2+3p^3+p^4$

$N(p,p,p^2,p^2)$	$7+5p+8p^2+9p^3+6p^4+p^5$
$N(p,p,p^2,p^2;1)$	1
$N(p,p,p^2,p^2;p)$	$1+p+p^2+p^3$
$N(p,p,p^2,p^2;p^2)$	$1+p+2p^2+2p^3+2p^4$
$N(p,p,p^2,p^2;p^3)$	$1+p+2p^2+3p^3+2p^4+p^5$

$N(p,p,p^2,p^3)$	$8+6p+10p^2+12p^3+10p^4+2p^5$
$N(p,p,p^2,p^3;1)$	1
$N(p,p,p^2,p^3;p)$	$1+p+p^2+p^3$
$N(p,p,p^2,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4$
$N(p,p,p^2,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+p^5$

$N(p,p,p^3,p^3)$	$9+7p+12p^2+15p^3+14p^4+6p^5+p^6$
$N(p,p,p^3,p^3;1)$	1
$N(p,p,p^3,p^3;p)$	$1+p+p^2+p^3$
$N(p,p,p^3,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4$
$N(p,p,p^3,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+2p^5$
$N(p,p,p^3,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+2p^5+p^6$

$N(p,p^2,p^2,p^2)$	$8+6p+10p^2+12p^3+10p^4+6p^5+2p^6$
$N(p,p^2,p^2,p^2;1)$	1
$N(p,p^2,p^2,p^2;p)$	$1+p+p^2+p^3$
$N(p,p^2,p^2,p^2;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5$
$N(p,p^2,p^2,p^2;p^3)$	$1+p+2p^2+3p^3+3p^4+2p^5+p^6$

$N(p,p^2,p^2,p^3)$	$9+7p+12p^2+15p^3+14p^4+11p^5+4p^6$
$N(p,p^2,p^2,p^3;1)$	1
$N(p,p^2,p^2,p^3;p)$	$1+p+p^2+p^3$
$N(p,p^2,p^2,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5$
$N(p,p^2,p^2,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+p^6$
$N(p,p^2,p^2,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+3p^5+2p^6$

$N(p,p^2,p^3,p^3)$	$10+8p+14p^2+18p^3+18p^4+16p^5+10p^6+2p^7$
$N(p,p^2,p^3,p^3;1)$	1
$N(p,p^2,p^3,p^3;p)$	$1+p+p^2+p^3$
$N(p,p^2,p^3,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5$
$N(p,p^2,p^3,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+2p^6$
$N(p,p^2,p^3,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+4p^5+3p^6+p^7$

$N(p,p^3,p^3,p^3)$	$11+9p+16p^2+21p^3+22p^4+21p^5+16p^6+9p^7+3p^8$
$N(p,p^3,p^3,p^3;1)$	1
$N(p,p^3,p^3,p^3;p)$	$1+p+p^2+p^3$
$N(p,p^3,p^3,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5$
$N(p,p^3,p^3,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+2p^6+p^7$
$N(p,p^3,p^3,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+4p^5+4p^6+2p^7+p^8$
$N(p,p^3,p^3,p^3;p^5)$	$1+p+2p^2+3p^3+4p^4+5p^5+4p^6+3p^7+p^8$

$N(p^2,p^2,p^2,p^2)$	$9+7p+12p^2+15p^3+14p^4+11p^5+9p^6+3p^7+p^8$
$N(p^2,p^2,p^2,p^2;1)$	1
$N(p^2,p^2,p^2,p^2;p)$	$1+p+p^2+p^3$
$N(p^2,p^2,p^2,p^2;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5+p^6$
$N(p^2,p^2,p^2,p^2;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+2p^6+p^7$
$N(p^2,p^2,p^2,p^2;p^4)$	$1+p+2p^2+3p^3+4p^4+3p^5+3p^6+p^7+p^8$

$N(p^2,p^2,p^2,p^3)$	$10+8p+14p^2+18p^3+18p^4+16p^5+16p^6+6p^7+2p^8$
$N(p^2,p^2,p^2,p^3;1)$	1
$N(p^2,p^2,p^2,p^3;p)$	$1+p+p^2+p^3$
$N(p^2,p^2,p^2,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5+p^6$
$N(p^2,p^2,p^2,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+3p^6+p^7$
$N(p^2,p^2,p^2,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+4p^5+4p^6+2p^7+p^8$

$N(p^2,p^2,p^3,p^3)$	$11+9p+16p^2+21p^3+22p^4+21p^5+23p^6+14p^7+6p^8+p^9$
$N(p^2,p^2,p^3,p^3;1)$	1
$N(p^2,p^2,p^3,p^3;p)$	$1+p+p^2+p^3$
$N(p^2,p^2,p^3,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5+p^6$
$N(p^2,p^2,p^3,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+3p^6+2p^7$
$N(p^2,p^2,p^3,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+4p^5+5p^6+3p^7+2p^8$
$N(p^2,p^2,p^3,p^3;p^5)$	$1+p+2p^2+3p^3+4p^4+5p^5+5p^6+4p^7+2p^8+p^9$

$N(p^2,p^3,p^3,p^3)$	$12+10p+18p^2+24p^3+26p^4+26p^5+30p^6+22p^7+16p^8+6p^9+2p^{10}$
$N(p^2,p^3,p^3,p^3;1)$	1
$N(p^2,p^3,p^3,p^3;p)$	$1+p+p^2+p^3$
$N(p^2,p^3,p^3,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5+p^6$
$N(p^2,p^3,p^3,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+3p^6+2p^7+p^8$
$N(p^2,p^3,p^3,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+4p^5+5p^6+4p^7+3p^8+p^9$
$N(p^2,p^3,p^3,p^3;p^5)$	$1+p+2p^2+3p^3+4p^4+5p^5+6p^6+5p^7+4p^8+2p^9+p^{10}$

$N(p^3,p^3,p^3,p^3)$	$13+11p+20p^2+27p^3+30p^4+31p^5+37p^6+30p^7+26p^8+18p^9$ $+9p^{10}+3p^{11}+p^{12}$
$N(p^3,p^3,p^3,p^3;1)$	1
$N(p^3,p^3,p^3,p^3;p)$	$1+p+p^2+p^3$
$N(p^3,p^3,p^3,p^3;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5+p^6$
$N(p^3,p^3,p^3,p^3;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+3p^6+2p^7+p^8+p^9$
$N(p^3,p^3,p^3,p^3;p^4)$	$1+p+2p^2+3p^3+4p^4+4p^5+5p^6+4p^7+4p^8+2p^9+p^{10}$
$N(p^3,p^3,p^3,p^3;p^5)$	$1+p+2p^2+3p^3+4p^4+5p^5+6p^6+6p^7+5p^8+4p^9+2p^{10}+p^{11}$
$N(p^3,p^3,p^3,p^3;p^6)$	$1+p+2p^2+3p^3+4p^4+5p^5+7p^6+6p^7+6p^8+4p^9+3p^{10}+p^{11}+p^{12}$

$N(p^4,p^4,p^4,p^4)$	$17+15p+28p^2+39p^3+46p^4+51p^5+65p^6+62p^7+66p^8+66p^9$ $+58p^{10}+46p^{11}+35p^{12}+18p^{13}+9p^{14}+3p^{15}+p^{16}$
$N(p^4,p^4,p^4,p^4;1)$	1
$N(p^4,p^4,p^4,p^4;p)$	$1+p+p^2+p^3$
$N(p^4,p^4,p^4,p^4;p^2)$	$1+p+2p^2+2p^3+2p^4+p^5+p^6$
$N(p^4,p^4,p^4,p^4;p^3)$	$1+p+2p^2+3p^3+3p^4+3p^5+3p^6+2p^7+p^8+p^9$
$N(p^4,p^4,p^4,p^4;p^4)$	$1+p+2p^2+3p^3+4p^4+4p^5+5p^6+4p^7+4p^8+3p^9+2p^{10}+p^{11}+p^{12}$
$N(p^4,p^4,p^4,p^4;p^5)$	$1+p+2p^2+3p^3+4p^4+5p^5+6p^6+6p^7+6p^8+6p^9+5p^{10}+4p^{11}+2p^{12}$ $+p^{13}$
$N(p^4,p^4,p^4,p^4;p^6)$	$1+p+2p^2+3p^3+4p^4+5p^5+7p^6+7p^7+8p^8+8p^9+8p^{10}+6p^{11}+5p^{12}$ $+2p^{13}+p^{14}$
$N(p^4,p^4,p^4,p^4;p^7)$	$1+p+2p^2+3p^3+4p^4+5p^5+7p^6+8p^7+9p^8+10p^9+9p^{10}+8p^{11}+6p^{12}$ $+4p^{13}+2p^{14}+p^{15}$
$N(p^4,p^4,p^4,p^4;p^8)$	$1+p+2p^2+3p^3+4p^4+5p^5+7p^6+8p^7+10p^8+10p^9+10p^{10}+8p^{11}+7p^{12}$ $+4p^{13}+3p^{14}+p^{15}+p^{16}$

5.2 Values of $N(n)$

The values of $N(n)$ for $1 \leqslant n \leqslant 30$ are given by the next table.

n	$N(n)$	n	$N(n)$	n	$N(n)$
1	1	11	19 156	21	774 224
2	67	12	420 396	22	1 283 452
3	212	13	35 872	23	318 532
4	1 983	14	244 684	24	9 187 868
5	1 120	15	237 440	25	810 969
6	14 204	16	821 335	26	2 403 424
7	3 652	17	99 472	27	2 222 704
8	43 339	18	1 610 211	28	7 241 916
9	24 033	19	152 404	29	783 904
10	75 040	20	2 220 960	30	15 908 480

6 Acknowledgment

The author thanks Mátyás Koniorczyk for his assistance during computations.

References

[1] K. Bauer, D. Sen, P. Zvengrowski, A generalized Goursat lemma, *Tatra Mt. Math. Publ.* **64** (2015), 1–19.

[2] G. Bhowmik, Evaluation of divisor functions of matrices, *Acta Arith.* **74** (1996), 155–159.

[3] L. M. Butler, A unimodality result in the enumeration of subgroups of a finite abelian group, *Proc. Amer. Math. Soc.* **101** (1987), 771–775.

[4] L. M. Butler, *Subgroup Lattices and Symmetric Functions*, Mem. Amer. Math. Soc., vol. 112, no. **539**, 1994.

[5] G. Călugăreanu, The total number of subgroups of a finite abelian group, *Sci. Math. Jpn.* **60** (2004), 157–167.

[6] C. Y. Chew, A. Y. M. Chin, and C. S. Lim, The number of subgroups of a finite abelian p-group of rank 4, AIP Conference Proceedings 1682, 040002 (2015). http://dx.doi.org/10.1063/1.4932475

[7] M. Golasiński and D. L. Gonçalves, On automorphisms of finite abelian p-groups, *Math. Slovaca* **58** (2008), 405–412.

[8] M. Hampejs, N. Holighaus, L. Tóth, and C. Wiesmeyr, Representing and counting the subgroups of the group $\mathbb{Z}_m \times \mathbb{Z}_n$, *Journal of Numbers*, vol. 2014, Article ID 491428.

[9] M. Hampejs and L. Tóth, On the subgroups of finite Abelian groups of rank three, *Annales Univ. Sci. Budapest., Sect. Comp.* **39** (2013), 111–124.

[10] Ch. J. Hillar and D. L. Rhea, Automorphisms of finite abelian groups, *Amer. Math. Monthly* **114** (2007), 917–923.

[11] Ju-Mok Oh, An explicit formula for the number of subgroups of a finite abelian p-group up to rank 3, *Commun. Korean Math. Soc.* **28** (2013), 649–667.

[12] J. Petrillo, Counting subgroups in a direct product of finite cyclic groups, *College Math. J.* **42** (2011), 215–222.

[13] V. N. Shokuev, An expression for the number of subgroups of a given order of a finite p-group, *Math. Notes* (Translation of Mat. Zametki) **12** (1972), 774–778.

[14] T. Stehling, On computing the number of subgroups of a finite Abelian group, *Combinatorica* **12** (1992), 475–479.

[15] Y. Takegahara, On Butler's unimodality result, *Combinatorica* **18** (1998), 437–439.

[16] M. Tărnăuceanu, An arithmetic method of counting the subgroups of a finite abelian group, *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)* **53(101)** (2010), 373–386.

[17] M. Tărnăuceanu and L. Tóth, On the number of subgroups of a given exponent in a finite abelian group, *Publ. Inst. Math. (Beograd) (N.S.)*, accepted. http://arxiv.org/abs/1507.00532

[18] L. Tóth, Subgroups of finite Abelian groups having rank two via Goursat's lemma, *Tatra Mt. Math. Publ.* **59** (2014), 93–103. http://arxiv.org/abs/1312.1485

[19] L. Tóth, Multiplicative arithmetic functions of several variables: a survey, in *Mathematics Without Boundaries, Surveys in Pure Mathematics*. Th. M. Rassias, P. Pardalos (Eds.), Springer, New York, 2014, 483–514. http://arxiv.org/abs/1310.7053

László Tóth
Department of Mathematics
University of Pécs
Ifjúság útja 6, H-7624 Pécs, Hungary
E-mail: ltoth@gamma.ttk.pte.hu